

Power 18% Goat Starter

MEDICATED

A Complete Feed for Goats.

For the prevention of coccidiosis caused by *Eimeria crandalis*, *Eimeria christenseni* and *Eimeria ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) 20.0 g/ton

GUARANTEED ANALYSIS

Crude Protein	(Min)		18.0 %
* (This includes not more than 1.70 % equivalent non-protein nitrogen)			
Crude Fat	(Min)		5.5 %
Crude Fiber	(Max)		9.0 %
Acid Detergent Fiber	(Max)		11.0 %
Calcium	(Min)	0.4	(Max) 0.8 %
Phosphorus	(Min)		0.4 %
Salt	(Min)	0.3	(Max) 0.6 %
Zinc	(Min)		150.0 ppm
Copper	(Min)		55.0 ppm
Selenium	(Min)		0.6 ppm
Vitamin A	(Min)		70,000.0 IU/lb
Vitamin E	(Min)		70.0 IU/lb

LIST OF INGREDIENTS

Corn, Soybean Meal, Corn Distillers Dried Grains with Solubles, Molasses Products, Soybean Hulls, Wheat Middlings, Beet Pulp, Cottonseed Hulls, Soybean Oil, Calcium Carbonate, Ammonium Chloride, Salt, Dried Saccharomyces Cerevisiae Fermentation Product, Artificial Flavor, Thiamin Mononitrate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydrodioxide, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid.

FEEDING DIRECTIONS

Feed at a rate of 2.5 to 4.0% of animal body weight to reach desired body condition. Typical consumption is 2 lbs of grain mix. Provide free choice good quality alfalfa hay and/or grass hay. Provide an abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further feeding directions consult your supplier or nutritionist.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats. CAUTION: This feed contains added copper, do not feed to sheep. CAUTION: Do not exceed 0.3 mg/kg of selenium in the complete feed.

Manufactured By:

Perry Milling Feed & Supply LLC
102 West Front Street
Perry, KS 66073
(785) 597-5122

No warranty of results is made due to differences in environment, genetics, health, and management. This product was made in a feed manufacturing facility that does not handle or store products containing animal protein prohibited in ruminant feed.
g180r2pk power 18% goat starter (180622) 3/29/2019

NET WT 50 LB (22.68 kg) or BULK

Code No.: G180R2PK

MEDICATED

Power 18% Goat Starter

MEDICATED

A Complete Feed for Goats.

For the prevention of coccidiosis caused by *Eimeria crandalis*, *Eimeria christenseni* and *Eimeria ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) 20.0 g/ton

GUARANTEED ANALYSIS

Crude Protein	(Min)		18.0 %
* (This includes not more than 1.70 % equivalent non-protein nitrogen)			
Crude Fat	(Min)		5.5 %
Crude Fiber	(Max)		9.0 %
Acid Detergent Fiber	(Max)		11.0 %
Calcium	(Min)	0.4	(Max) 0.8 %
Phosphorus	(Min)		0.4 %
Salt	(Min)	0.3	(Max) 0.6 %
Zinc	(Min)		150.0 ppm
Copper	(Min)		55.0 ppm
Selenium	(Min)		0.6 ppm
Vitamin A	(Min)		70,000.0 IU/lb
Vitamin E	(Min)		70.0 IU/lb

LIST OF INGREDIENTS

Corn, Soybean Meal, Corn Distillers Dried Grains with Solubles, Molasses Products, Soybean Hulls, Wheat Middlings, Beet Pulp, Cottonseed Hulls, Soybean Oil, Calcium Carbonate, Ammonium Chloride, Salt, Dried Saccharomyces Cerevisiae Fermentation Product, Artificial Flavor, Thiamin Mononitrate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydrodioxide, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid.

FEEDING DIRECTIONS

Feed at a rate of 2.5 to 4.0% of animal body weight to reach desired body condition. Typical consumption is 2 lbs of grain mix. Provide free choice good quality alfalfa hay and/or grass hay. Provide an abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further feeding directions consult your supplier or nutritionist.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats. CAUTION: This feed contains added copper, do not feed to sheep. CAUTION: Do not exceed 0.3 mg/kg of selenium in the complete feed.

Manufactured By:

Perry Milling Feed & Supply LLC
102 West Front Street
Perry, KS 66073
(785) 597-5122

No warranty of results is made due to differences in environment, genetics, health, and management. This product was made in a feed manufacturing facility that does not handle or store products containing animal protein prohibited in ruminant feed.
g180r2pk power 18% goat starter (180622) 3/29/2019

NET WT 50 LB (22.68 kg) or BULK

Code No.: G180R2PK

MEDICATED

Power 18% Goat Starter

MEDICATED

A Complete Feed for Goats.

For the prevention of coccidiosis caused by *Eimeria crandalis*, *Eimeria christenseni* and *Eimeria ninakohlyakimovae*.

ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) 20.0 g/ton

GUARANTEED ANALYSIS

Crude Protein	(Min)		18.0 %
* (This includes not more than 1.70 % equivalent non-protein nitrogen)			
Crude Fat	(Min)		5.5 %
Crude Fiber	(Max)		9.0 %
Acid Detergent Fiber	(Max)		11.0 %
Calcium	(Min)	0.4	(Max) 0.8 %
Phosphorus	(Min)		0.4 %
Salt	(Min)	0.3	(Max) 0.6 %
Zinc	(Min)		150.0 ppm
Copper	(Min)		55.0 ppm
Selenium	(Min)		0.6 ppm
Vitamin A	(Min)		70,000.0 IU/lb
Vitamin E	(Min)		70.0 IU/lb

LIST OF INGREDIENTS

Corn, Soybean Meal, Corn Distillers Dried Grains with Solubles, Molasses Products, Soybean Hulls, Wheat Middlings, Beet Pulp, Cottonseed Hulls, Soybean Oil, Calcium Carbonate, Ammonium Chloride, Salt, Dried Saccharomyces Cerevisiae Fermentation Product, Artificial Flavor, Thiamin Mononitrate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Sodium Selenite, Ethylenediamine Dihydrodioxide, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid.

FEEDING DIRECTIONS

Feed at a rate of 2.5 to 4.0% of animal body weight to reach desired body condition. Typical consumption is 2 lbs of grain mix. Provide free choice good quality alfalfa hay and/or grass hay. Provide an abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further feeding directions consult your supplier or nutritionist.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats. CAUTION: This feed contains added copper, do not feed to sheep. CAUTION: Do not exceed 0.3 mg/kg of selenium in the complete feed.

Manufactured By:

Perry Milling Feed & Supply LLC
102 West Front Street
Perry, KS 66073
(785) 597-5122

No warranty of results is made due to differences in environment, genetics, health, and management. This product was made in a feed manufacturing facility that does not handle or store products containing animal protein prohibited in ruminant feed.
g180r2pk power 18% goat starter (180622) 3/29/2019

NET WT 50 LB (22.68 kg) or BULK

Code No.: G180R2PK

MEDICATED