



PERRY MILLING

EST. FEED & SUPPLY 1992 COMMENTED



Power 14% Goat Finisher

MEDICATED

A Complete Feed for Goats.

For the prevention of coccidiosis caused by Eimeria crandalis. Eimeria christenseni and Eimeria ninakohlyakimovae.

ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium)

GUAR	ANTEE	D ANAL	YSIS

G	JAKANII	EED A	NALISIS	
Crude Protein			(Min)	14.0 %
* (This includes not n	nore than 1.	70 % eq	uivalent non-	protein nitrogen)
Crude Fat			(Min)	5.5 %
Crude Fiber			(Max)	10.0 %
Acid Detergent Fiber			(Max)	10.0 %
Calcium	(Min)	0.3	(Max)	0.8 %
Phosphorus			(Min)	0.3 %
Salt	(Min)	0.4	(Max)	0.9 %
Zinc			(Min)	100.0 ppm
Copper			(Min)	10.0 ppm
Selenium			(Min)	0.3 ppm
Vitamin A			(Min)	30,000.0 IU/lb
Vitamin E			(Min)	30.0 IU/lb

LIST OF INGREDIENTS

Corn, Molasses Products, Wheat Middlings, Soybean Meal, Corn Distillers Dried Grains with Solubles, Soybean Hulls, Beet Pulp, Cottonseed Hulls, Soybean Oil, Calcium Carbonate, Ammonium Chloride, Salt, Potassium Chloride, Artificial Flavor, Dried Saccharomyces Cerevisiae Fermentation Product, Thiamin Mononitrate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid, Zinc Sulfate, Zinc Oxide, Ferrous Sulfate, Manganese Sulfate, Manganese Oxide, Ethylenediamine Dihydriodide, Calcium Iodate, Cobalt Sulfate, Cobalt Carbonate, Sodium Selenite, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Copper Sulfate, Silicon Dioxide, Ethoxyguin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid.

FEEDING DIRECTIONS

Feed at a rate of 2.5 to 4.5% of animal body weight to reach desired body condition. Typical consumption is 4 lbs of grain mix. Provide free choice good quality alfalfa hay and/or grass hay. Provide an abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further feeding directions consult your supplier or nutritionist.



Manufactured By:

Perry Milling Feed & Supply LLC 102 West Front Street Perrv. KS 66073 (785) 597-5122

No warranty of results is made due to differences in environment, genetics, health, and management. This product was made in a feed manufacturing facility that does not handle or store products containing animal protein prohibited in ruminant feed. 3/29/2019 g140r2pk power 14% goat finisher (180622)

NET WT 50 LB (22.68 kg) or BULK

Code No.: G140R2PK















PERRY MILLING

EST. FEED & SUPPLY



Power 14% Goat Finisher

MEDICATED

A Complete Feed for Goats.

For the prevention of coccidiosis caused by Eimeria crandalis. Eimeria christenseni and Eimeria ninakohlyakimovae.

ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) 20.0 a/ton

MOHENSIII (as MOHENS	in Souluini)			20.0 g/t011	
GUARANTEED ANALYSIS					
Crude Protein			(Min)	14.0 %	
* (This includes not more than 1.70 % equivalent non-protein nitrogen)					
Crude Fat			(Min)	5.5 %	
Crude Fiber			(Max)	10.0 %	
Acid Detergent Fiber			(Max)	10.0 %	
Calcium	(Min)	0.3	(Max)	0.8 %	
Phosphorus			(Min)	0.3 %	
Salt	(Min)	0.4	(Max)	0.9 %	
Zinc			(Min)	100.0 ppm	
Copper			(Min)	10.0 ppm	
Selenium			(Min)	0.3 ppm	
Vitamin A			(Min)	30,000.0 IU/lb	
Vitamin E			(Min)	30.0 IU/lb	

LIST OF INGREDIENTS

Corn, Molasses Products, Wheat Middlings, Sovbean Meal, Corn Distillers Dried Grains with Solubles, Soybean Hulls, Beet Pulp, Cottonseed Hulls, Soybean Oil, Calcium Carbonate, Ammonium Chloride, Salt, Potassium Chloride, Artificial Flavor, Dried Saccharomyces Cerevisiae Fermentation Product, Thiamin Mononitrate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid, Zinc Sulfate, Zinc Oxide, Ferrous Sulfate, Manganese Sulfate, Manganese Oxide, Ethylenediamine Dihydriodide, Calcium Iodate, Cobalt Sulfate, Cobalt Carbonate, Sodium Selenite, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Copper Sulfate, Silicon Dioxide, Ethoxyguin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a

FEEDING DIRECTIONS

Feed at a rate of 2.5 to 4.5% of animal body weight to reach desired body condition. Typical consumption is 4 lbs of grain mix. Provide free choice good quality alfalfa hay and/or grass hay. Provide an abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further feeding directions consult your supplier or nutritionist.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding

directions, as reduced average daily gains may result. Do not feed to lactating goats.

CAUTION: This feed contains added copper, do not feed to sheep, CAUTION: Do not exceed 0.3 mg/kg of selenium in the complete feed.

Code No.: G140R2PK

Manufactured By:

Perry Milling Feed & Supply LLC 102 West Front Street Perrv. KS 66073 (785) 597-5122

No warranty of results is made due to differences in environment, genetics, health, and management. This product was made in a feed manufacturing facility that does not handle or store products containing animal protein prohibited in ruminant feed. 3/29/2019 g140r2pk power 14% goat finisher (180622)

NET WT 50 LB (22.68 kg) or BULK



Power 14% Goat Finisher

MEDICATED

A Complete Feed for Goats.

For the prevention of coccidiosis caused by Eimeria crandalis. Eimeria christenseni and Eimeria ninakohlyakimovae.

ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) 20.0 a/ton

GUARANTEED ANALYSIS

Crude Protein 14 0 % * (This includes not more than 1.70 % equivalent non-protein nitrogen)

Crude Fat			(Min)	5.5 %
Crude Fiber			(Max)	10.0 %
Acid Detergent Fiber			(Max)	10.0 %
Calcium	(Min)	0.3	(Max)	0.8 %
Phosphorus			(Min)	0.3 %
Salt	(Min)	0.4	(Max)	0.9 %
Zinc			(Min)	100.0 ppm
Copper			(Min)	10.0 ppm
Selenium			(Min)	0.3 ppm
Vitamin A			(Min)	30,000.0 IU/lb
Vitamin E			(Min)	30.0 IU/lb

LIST OF INGREDIENTS

Corn, Molasses Products, Wheat Middlings, Sovbean Meal, Corn Distillers Dried Grains with Solubles, Soybean Hulls, Beet Pulp, Cottonseed Hulls, Soybean Oil, Calcium Carbonate, Ammonium Chloride, Salt, Potassium Chloride, Artificial Flavor, Dried Saccharomyces Cerevisiae Fermentation Product, Thiamin Mononitrate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid, Zinc Sulfate, Zinc Oxide, Ferrous Sulfate, Manganese Sulfate, Manganese Oxide, Ethylenediamine Dihydriodide, Calcium Iodate, Cobalt Sulfate, Cobalt Carbonate, Sodium Selenite, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Copper Sulfate, Silicon Dioxide, Ethoxyguin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid.

FEEDING DIRECTIONS

Feed at a rate of 2.5 to 4.5% of animal body weight to reach desired body condition. Typical consumption is 4 lbs of grain mix. Provide free choice good quality alfalfa hay and/or grass hay. Provide an abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further feeding directions consult your supplier or nutritionist.



CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats. CAUTION: This feed contains added copper, do not feed to sheep, CAUTION: Do not exceed 0.3 mg/kg of selenium in the complete feed.



Perry Milling Feed & Supply LLC 102 West Front Street Perry, KS 66073 (785) 597-5122

No warranty of results is made due to differences in environment, genetics, health, and management. This product was made in a feed manufacturing facility that does not handle or store products containing animal protein prohibited in ruminant feed. 3/29/2019 g140r2pk power 14% goat finisher (180622)

NET WT 50 LB (22.68 kg) or BULK

Code No.: G140R2PK

MEDICATED