

# Performance

## 16% Exacta 150 Creep

**MEDICATED**

A Complete Feed for Beef Cattle.

For improved feed efficiency.

**ACTIVE DRUG INGREDIENT**

Monensin (As Monensin Sodium) ..... 75.0 g/ton

**GUARANTEED ANALYSIS**

Crude Protein	(Min)		16.0 %
Crude Fat	(Min)		4.0 %
Crude Fiber	(Max)		8.0 %
Calcium	(Min)	0.9	(Max) 1.4 %
Phosphorus	(Min)		0.6 %
Salt	(Min)	0.4	(Max) 0.8 %
Zinc	(Min)		180.0 ppm
Copper	(Min)		50.0 ppm
Selenium	(Min)		0.6 ppm
Vitamin A	(Min)		15,000.0 IU/lb
Vitamin E	(Min)		15.0 IU/lb

**LIST OF INGREDIENTS**

Wheat Middlings, Corn, Corn Gluten Feed, Corn Distillers Dried Grains with Solubles, Molasses Products, Calcium Carbonate, Calcium Chloride, Salt, Potassium Chloride, Thiamin Mononitrate, Sodium Selenite, Copper Sulfate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid, Zinc Sulfate, Zinc Oxide, Ferrous Sulfate, Manganese Sulfate, Manganese Oxide, Ethylenediamine Dihydrochloride, Calcium Iodate, Cobalt Sulfate, Cobalt Carbonate.

**FEEDING DIRECTIONS**

Feed as a complete feed, along with free-choice, good quality grass/alfalfa hay or feed on pasture. Feed at a rate of 5 to 10 lb/head/day of grain mix to provide 187.5 to 375 mg/head/day of Monensin (Rumensin). Cattle should consume approximately 1.5% of their bodyweight in grain mix. Provide an abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further directions consult your supplier or nutritionist.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. CAUTION: This feed contains added copper, do not feed to sheep. CAUTION: Do not exceed 0.3 mg/kg of selenium in the complete feed.

**Manufactured By:**

Perry Milling  
102 West Front Street  
Perry, KS 66073  
(785) 597-5122

No warranty of results is made due to differences in environment, genetics, health, and management. This product was made in a feed manufacturing facility that does not handle or store products containing animal protein prohibited in ruminant feed.

a160r1pk performance 16% exacta 150 creep (180406).docx 1/25/2019  
NET WT 50 LB (22.68 kg) or BULK

**Code No.: A160R1PK**

**MEDICATED**

# Performance

## 16% Exacta 150 Creep

**MEDICATED**

A Complete Feed for Beef Cattle.

For improved feed efficiency.

**ACTIVE DRUG INGREDIENT**

Monensin (As Monensin Sodium) ..... 75.0 g/ton

**GUARANTEED ANALYSIS**

Crude Protein	(Min)		16.0 %
Crude Fat	(Min)		4.0 %
Crude Fiber	(Max)		8.0 %
Calcium	(Min)	0.9	(Max) 1.4 %
Phosphorus	(Min)		0.6 %
Salt	(Min)	0.4	(Max) 0.8 %
Zinc	(Min)		180.0 ppm
Copper	(Min)		50.0 ppm
Selenium	(Min)		0.6 ppm
Vitamin A	(Min)		15,000.0 IU/lb
Vitamin E	(Min)		15.0 IU/lb

**LIST OF INGREDIENTS**

Wheat Middlings, Corn, Corn Gluten Feed, Corn Distillers Dried Grains with Solubles, Molasses Products, Calcium Carbonate, Calcium Chloride, Salt, Potassium Chloride, Thiamin Mononitrate, Sodium Selenite, Copper Sulfate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid, Zinc Sulfate, Zinc Oxide, Ferrous Sulfate, Manganese Sulfate, Manganese Oxide, Ethylenediamine Dihydrochloride, Calcium Iodate, Cobalt Sulfate, Cobalt Carbonate.

**FEEDING DIRECTIONS**

Feed as a complete feed, along with free-choice, good quality grass/alfalfa hay or feed on pasture. Feed at a rate of 5 to 10 lb/head/day of grain mix to provide 187.5 to 375 mg/head/day of Monensin (Rumensin). Cattle should consume approximately 1.5% of their bodyweight in grain mix. Provide an abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further directions consult your supplier or nutritionist.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. CAUTION: This feed contains added copper, do not feed to sheep. CAUTION: Do not exceed 0.3 mg/kg of selenium in the complete feed.

**Manufactured By:**

Perry Milling  
102 West Front Street  
Perry, KS 66073  
(785) 597-5122

No warranty of results is made due to differences in environment, genetics, health, and management. This product was made in a feed manufacturing facility that does not handle or store products containing animal protein prohibited in ruminant feed.

a160r1pk performance 16% exacta 150 creep (180406).docx 1/25/2019  
NET WT 50 LB (22.68 kg) or BULK

**Code No.: A160R1PK**

**MEDICATED**

# Performance

## 16% Exacta 150 Creep

**MEDICATED**

A Complete Feed for Beef Cattle.

For improved feed efficiency.

**ACTIVE DRUG INGREDIENT**

Monensin (As Monensin Sodium) ..... 75.0 g/ton

**GUARANTEED ANALYSIS**

Crude Protein	(Min)		16.0 %
Crude Fat	(Min)		4.0 %
Crude Fiber	(Max)		8.0 %
Calcium	(Min)	0.9	(Max) 1.4 %
Phosphorus	(Min)		0.6 %
Salt	(Min)	0.4	(Max) 0.8 %
Zinc	(Min)		180.0 ppm
Copper	(Min)		50.0 ppm
Selenium	(Min)		0.6 ppm
Vitamin A	(Min)		15,000.0 IU/lb
Vitamin E	(Min)		15.0 IU/lb

**LIST OF INGREDIENTS**

Wheat Middlings, Corn, Corn Gluten Feed, Corn Distillers Dried Grains with Solubles, Molasses Products, Calcium Carbonate, Calcium Chloride, Salt, Potassium Chloride, Thiamin Mononitrate, Sodium Selenite, Copper Sulfate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid, Zinc Sulfate, Zinc Oxide, Ferrous Sulfate, Manganese Sulfate, Manganese Oxide, Ethylenediamine Dihydrochloride, Calcium Iodate, Cobalt Sulfate, Cobalt Carbonate.

**FEEDING DIRECTIONS**

Feed as a complete feed, along with free-choice, good quality grass/alfalfa hay or feed on pasture. Feed at a rate of 5 to 10 lb/head/day of grain mix to provide 187.5 to 375 mg/head/day of Monensin (Rumensin). Cattle should consume approximately 1.5% of their bodyweight in grain mix. Provide an abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further directions consult your supplier or nutritionist.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and the amount of refusals fed should be taken into consideration to prevent Monensin overdosing. WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. CAUTION: This feed contains added copper, do not feed to sheep. CAUTION: Do not exceed 0.3 mg/kg of selenium in the complete feed.

**Manufactured By:**

Perry Milling  
102 West Front Street  
Perry, KS 66073  
(785) 597-5122

No warranty of results is made due to differences in environment, genetics, health, and management. This product was made in a feed manufacturing facility that does not handle or store products containing animal protein prohibited in ruminant feed.

a160r1pk performance 16% exacta 150 creep (180406).docx 1/25/2019  
NET WT 50 LB (22.68 kg) or BULK

**Code No.: A160R1PK**

**MEDICATED**