PERRY MILLING GOAT STARTER 19% MON25

MEDICATED

For the prevention of coccidiosis caused by *Eimeria crandallis*, *Eimeria christenseni* and *Eimeria ninakohlyakimovae* in goats maintained in confinement.

ACTIVE DRUG INGREDIENTS

	<u> </u>	<u> </u>		<u></u>			
Monensin			2	25		G	/TON
	GUAR	ANTE	D ANAL	YSIS			
Crude Protein				(Min)		19	9.0 %
This includes not	more tha	n 1.4%	equivalent	protein	from	non-p	orotein
Crude Fat				(Min)		3	3.6 %
Crude Fiber				(Max)		8	3.0 %
Acid Detergent				(Max)		é	9.0 %
Calcium	(1	Min)	1	(Max)		•	1.5 %
Phosphorus				(Min)		(0.3 %
Salt	(1	Min)	0.1	(Max)		(0.6 %
Copper	(Min)	25	(Min)		75	ppm
Selenium				(Min)		.375	ppm
Vitamin A				(Min)	4,	300	IU/lb

INGREDIENTS

Grain products, plant protein products, processed grain by-products, molasses products, calcium carbonate, vegetable oil, animal protein products, ammonium chloride, salt, urea, natural flavor, artificial flavor, selenium yeast, hydrolyzed yeast, zinc proteinate, manganese proteinate, vitamin E supplement, calcium hydroxide, copper proteinate, sodium carboxymethylcellulose, roughage products, magnesium oxide, cobalt proteinate, ethylenediamine dihydroiodide, vitamin A supplement, thiamine mononitrate, vitamin D3 supplement, mineral oil

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper. This product has been formulated specifically for goats and is not intended for other species.

CAUTION: USE AS DIRECTED CAUTION: Follow label directions. The addition to feed of higher levels of this supplement containing added selenium is not permitted.

Feeding Directions: Perry Milling Goat Starter 19% MON25 is a supplement for starting goats. Thoroughly mix according to the table below to provide a mixture with 20 grams per ton of Monensin. Feed this mixture continuously as the sole ration. A source of good quality hay and a constant supply of fresh, clean water is also required.

	Starter Mix
Perry Milling Goat Starter 19% MON25, lbs	1,600
Non-medicated Feedstuff, lbs	400
Total lhs	2.000

WARNING: Do not feed to lactating goats.

CAUTION: Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Do not feed undiluted. Monensin medicated cattle and goat feeds are safe for cattle and goats only. Consumption by unapproved species may result in toxic reactions. Must be thoroughly mixed in feed before use. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats.

Net Weight 50 lb (22.68 kg) or bulk invoice.

PERRY MILLING GOAT STARTER 19% MON25

MEDICATED

For the prevention of coccidiosis caused by *Eimeria crandallis*, *Eimeria christenseni* and *Eimeria ninakohlyakimovae* in goats maintained in confinement.

ACTIVE DRUG INGREDIENTS

ACTIVE DRUG INGREDIENTS					
Monensin		2	25	G/TON	
	GUARANTEE	D ANAL	YSIS		
Crude Protein			(Min)	19.0 %	
This includes not	more than 1.4%	equivalent	protein	from non-protein	
Crude Fat			(Min)	3.6 %	
Crude Fiber			(Max)	8.0 %	
Acid Detergent			(Max)	9.0 %	
Calcium	(Min)	1	(Max)	1.5 %	
Phosphorus			(Min)	0.3 %	
Salt	(Min)	0.1	(Max)	0.6 %	
Copper	(Min)	25	(Min)	75 ppm	
Selenium			(Min)	.375 ppm	
Vitamin A			(Min)	4.300 IU/lb	

INGREDIENTS

Grain products, plant protein products, processed grain by-products, molasses products, calcium carbonate, vegetable oil, animal protein products, ammonium chloride, salt, urea, natural flavor, artificial flavor, selenium yeast, hydrolyzed yeast, zinc proteinate, manganese proteinate, vitamin E supplement, calcium hydroxide, copper proteinate, sodium carboxymethylcellulose, roughage products, magnesium oxide, cobalt proteinate, ethylenediamine dihydroiodide, vitamin A supplement, thiamine mononitrate, vitamin D3 supplement, mineral oil

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper. This product has been formulated specifically for goats and is not intended for other species.

CAUTION: USE AS DIRECTED CAUTION: Follow label directions. The addition to feed of higher levels of this supplement containing added selenium is not permitted.

Feeding Directions: Perry Milling Goat Starter 19% MON25 is a supplement for starting goats. Thoroughly mix according to the table below to provide a mixture with 20 grams per ton of Monensin. Feed this mixture continuously as the sole ration. A source of good quality hay and a constant supply of fresh, clean water is also required.

	Starter Mix
Perry Milling Goat Starter 19% MON25, lbs	1,600
Non-medicated Feedstuff, lbs	400
Total lhs	2 000

WARNING: Do not feed to lactating goats.

CAUTION: Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Do not feed undiluted. Monensin medicated cattle and goat feeds are safe for cattle and goats only. Consumption by unapproved species may result in toxic reactions. Must be thoroughly mixed in feed before use. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats.

Net Weight 50 lb (22.68 kg) or bulk invoice.

PERRY MILLING GOAT STARTER 19% MON25

MEDICATED

For the prevention of coccidiosis caused by *Eimeria crandallis*, *Eimeria christenseni* and *Eimeria ninakohlyakimovae* in goats maintained in confinement.

ACTIVE DRUG INGREDIENTS

ACTIVE DRUG INGREDIENTS					
Monensin			25		
GUARANTEED ANALYSIS					
			(Min)		19.0 %
more than '	1.4%	equivalent	protein	from	non-protein
			(Min)		3.6 %
			(Max)		8.0 %
			(Max)		9.0 %
(Mir	า)	1	(Max)		1.5 %
			(Min)		0.3 %
(Mir	า)	0.1	(Max)		0.6 %
(Mi	in)	25	(Min)		75 ppm
,	•		(Min)		.375 ppm
			(Min)	4,	300 IU/lb
	GUARAN more than (Min	GUARANTEE	GUARANTEED ANAL more than 1.4% equivalent (Min) 1 (Min) 0.1	25 GUARANTEED ANALYSIS (Min) (Min) (Min) (Min) (Max) (Max) (Max) (Max) (Max) (Min) (Min) (Min) (Min) (Min) (Min) (Min) (25 (Min) (Min) (Min) (Min) (Min)	25 GUARANTEED ANALYSIS (Min) more than 1.4% equivalent protein from (Min) (Max) (Max) (Max) (Min) 1 (Max) (Min) (Min) (Min) 25 (Min) (Min) (Min)

INGREDIENTS

Grain products, plant protein products, processed grain by-products, molasses products, calcium carbonate, vegetable oil, animal protein products, ammonium chloride, salt, urea, natural flavor, artificial flavor, selenium yeast, hydrolyzed yeast, zinc proteinate, manganese proteinate, vitamin E supplement, calcium hydroxide, copper proteinate, sodium carboxymethylcellulose, roughage products, magnesium oxide, cobalt proteinate, ethylenediamine dihydroiodide, vitamin A supplement, thiamine mononitrate, vitamin D3 supplement, mineral oil

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper. This product has been formulated specifically for goats and is not intended for other species.

CAUTION: USE AS DIRECTED CAUTION: Follow label directions. The addition to feed of higher levels of this supplement containing added selenium is not permitted.

Feeding Directions: Perry Milling Goat Starter 19% MON25 is a supplement for starting goats. Thoroughly mix according to the table below to provide a mixture with 20 grams per ton of Monensin. Feed this mixture continuously as the sole ration. A source of good quality hay and a constant supply of fresh, clean water is also required.

	Starter Mix
Perry Milling Goat Starter 19% MON25, lbs	1,600
Non-medicated Feedstuff, lbs	400
Total, lbs	2,000

WARNING: Do not feed to lactating goats.

CAUTION: Do not allow horses or other equines access to formulations containing Monensin. Ingestion of Monensin by equines has been fatal. Do not feed undituted. Monensin medicated cattle and goat feeds are safe for cattle and goats only. Consumption by unapproved species may result in toxic reactions. Must be thoroughly mixed in feed before use. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Feeding undituted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats.

Net Weight 50 lb (22.68 kg) or bulk invoice.