

# ECS 20-20 AM with BOV Celmanax® and Selenium Yeast CALF MILK REPLACER MEDICATED

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in replacement calves. Feed during periods of coccidiosis exposure or when experience indicates coccidiosis is likely to be a problem.

## ACTIVE DRUG INGREDIENT

Lasalocid (as lasalocid sodium) . . . . . 90 g/ton

**WARNING:** Withdrawal period has not been established for lasalocid in pre-ruminating calves. Do not use in calves to be processed for veal.

**CAUTION:** Do not allow horses or other equines access to feeds containing lasalocid as ingestion may be fatal.

## GUARANTEED ANALYSIS

Crude Protein, Min. . . . . 20.00%  
Crude Fat, Min. . . . . 20.00%  
Crude Fiber, Max. . . . . 0.15%  
Calcium, Min. . . . . 0.75%  
Calcium, Max. . . . . 1.25%  
Phosphorus, Min. . . . . 0.70%  
Vitamin A, Min. . . . . 35,000 IU/lb  
Vitamin D<sub>3</sub>, Min. . . . . 7,500 IU/lb  
Vitamin E, Min. . . . . 150 IU/lb

## INGREDIENTS

Dried Whey, Dried Whey Protein Concentrate, Dried Whey Product, Animal and Vegetable Fat (Preserved with BHA and BHT), Dried Skimmed Milk, Lecithin, Dicalcium Phosphate, Calcium Carbonate, Yeast Culture, Yeast Extract, Hydrolyzed Yeast, L-Lysine Monohydrochloride, DL-Methionine, Vitamin A Supplement, Vitamin D<sub>3</sub> Supplement, Vitamin E Supplement, Ascorbic Acid, Magnesium Oxide, Zinc Sulfate, Ferrous Sulfate, Niacin Supplement, Manganese Sulfate, Calcium Pantothenate, Vitamin B<sub>12</sub> Supplement, Thiamine Mononitrate, Riboflavin Supplement, Biotin, Copper Sulfate, Pyridoxine Hydrochloride, Ethylenediamine Dihydriodide, Folic Acid, Choline Chloride, Cobalt Sulfate, Menadione Sodium Bisulfite Complex (Source of Vitamin K Activity), Selenium Yeast, Sodium Silico Aluminate, Mono and Diglycerides of Edible Fats or Oils, Natural and Artificial Flavors.

SEE FEEDING DIRECTIONS ON REVERSE SIDE

Net Weight 50 lb (22.68 kg)

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.

Manufactured for:

Eldon C. Stutsman, Inc.  
Hills, IA 52235 U.S.A.  
(319) 679-2281

MADE IN THE U.S.A.

488732

REV 1117

FI - 001

# ECS 20-20 AM with BOV Celmanax® and Selenium Yeast CALF MILK REPLACER MEDICATED

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in replacement calves. Feed during periods of coccidiosis exposure or when experience indicates coccidiosis is likely to be a problem.

## ACTIVE DRUG INGREDIENT

Lasalocid (as lasalocid sodium) . . . . . 90 g/ton

**WARNING:** Withdrawal period has not been established for lasalocid in pre-ruminating calves. Do not use in calves to be processed for veal.

**CAUTION:** Do not allow horses or other equines access to feeds containing lasalocid as ingestion may be fatal.

## GUARANTEED ANALYSIS

Crude Protein, Min. . . . . 20.00%  
Crude Fat, Min. . . . . 20.00%  
Crude Fiber, Max. . . . . 0.15%  
Calcium, Min. . . . . 0.75%  
Calcium, Max. . . . . 1.25%  
Phosphorus, Min. . . . . 0.70%  
Vitamin A, Min. . . . . 35,000 IU/lb  
Vitamin D<sub>3</sub>, Min. . . . . 7,500 IU/lb  
Vitamin E, Min. . . . . 150 IU/lb

## INGREDIENTS

Dried Whey, Dried Whey Protein Concentrate, Dried Whey Product, Animal and Vegetable Fat (Preserved with BHA and BHT), Dried Skimmed Milk, Lecithin, Dicalcium Phosphate, Calcium Carbonate, Yeast Culture, Yeast Extract, Hydrolyzed Yeast, L-Lysine Monohydrochloride, DL-Methionine, Vitamin A Supplement, Vitamin D<sub>3</sub> Supplement, Vitamin E Supplement, Ascorbic Acid, Magnesium Oxide, Zinc Sulfate, Ferrous Sulfate, Niacin Supplement, Manganese Sulfate, Calcium Pantothenate, Vitamin B<sub>12</sub> Supplement, Thiamine Mononitrate, Riboflavin Supplement, Biotin, Copper Sulfate, Pyridoxine Hydrochloride, Ethylenediamine Dihydriodide, Folic Acid, Choline Chloride, Cobalt Sulfate, Menadione Sodium Bisulfite Complex (Source of Vitamin K Activity), Selenium Yeast, Sodium Silico Aluminate, Mono and Diglycerides of Edible Fats or Oils, Natural and Artificial Flavors.

SEE FEEDING DIRECTIONS ON REVERSE SIDE

Net Weight 50 lb (22.68 kg)

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.

Manufactured for:

Eldon C. Stutsman, Inc.  
Hills, IA 52235 U.S.A.  
(319) 679-2281

MADE IN THE U.S.A.

488732

REV 1117

FI - 001

# ECS 20-20 AM with BOV Celmanax® and Selenium Yeast CALF MILK REPLACER MEDICATED

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in replacement calves. Feed during periods of coccidiosis exposure or when experience indicates coccidiosis is likely to be a problem.

## ACTIVE DRUG INGREDIENT

Lasalocid (as lasalocid sodium) . . . . . 90 g/ton

**WARNING:** Withdrawal period has not been established for lasalocid in pre-ruminating calves. Do not use in calves to be processed for veal.

**CAUTION:** Do not allow horses or other equines access to feeds containing lasalocid as ingestion may be fatal.

## GUARANTEED ANALYSIS

Crude Protein, Min. . . . . 20.00%  
Crude Fat, Min. . . . . 20.00%  
Crude Fiber, Max. . . . . 0.15%  
Calcium, Min. . . . . 0.75%  
Calcium, Max. . . . . 1.25%  
Phosphorus, Min. . . . . 0.70%  
Vitamin A, Min. . . . . 35,000 IU/lb  
Vitamin D<sub>3</sub>, Min. . . . . 7,500 IU/lb  
Vitamin E, Min. . . . . 150 IU/lb

## INGREDIENTS

Dried Whey, Dried Whey Protein Concentrate, Dried Whey Product, Animal and Vegetable Fat (Preserved with BHA and BHT), Dried Skimmed Milk, Lecithin, Dicalcium Phosphate, Calcium Carbonate, Yeast Culture, Yeast Extract, Hydrolyzed Yeast, L-Lysine Monohydrochloride, DL-Methionine, Vitamin A Supplement, Vitamin D<sub>3</sub> Supplement, Vitamin E Supplement, Ascorbic Acid, Magnesium Oxide, Zinc Sulfate, Ferrous Sulfate, Niacin Supplement, Manganese Sulfate, Calcium Pantothenate, Vitamin B<sub>12</sub> Supplement, Thiamine Mononitrate, Riboflavin Supplement, Biotin, Copper Sulfate, Pyridoxine Hydrochloride, Ethylenediamine Dihydriodide, Folic Acid, Choline Chloride, Cobalt Sulfate, Menadione Sodium Bisulfite Complex (Source of Vitamin K Activity), Selenium Yeast, Sodium Silico Aluminate, Mono and Diglycerides of Edible Fats or Oils, Natural and Artificial Flavors.

SEE FEEDING DIRECTIONS ON REVERSE SIDE

Net Weight 50 lb (22.68 kg)

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.

Manufactured for:

Eldon C. Stutsman, Inc.  
Hills, IA 52235 U.S.A.  
(319) 679-2281

MADE IN THE U.S.A.

488732

REV 1117

FI - 001

# ECS 20-20 AM with BOV Celmanax® and Selenium Yeast CALF MILK REPLACER MEDICATED

For the control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in replacement calves. Feed during periods of coccidiosis exposure or when experience indicates coccidiosis is likely to be a problem.

## ACTIVE DRUG INGREDIENT

Lasalocid (as lasalocid sodium) . . . . . 90 g/ton

**WARNING:** Withdrawal period has not been established for lasalocid in pre-ruminating calves. Do not use in calves to be processed for veal.

**CAUTION:** Do not allow horses or other equines access to feeds containing lasalocid as ingestion may be fatal.

## GUARANTEED ANALYSIS

Crude Protein, Min. . . . . 20.00%  
Crude Fat, Min. . . . . 20.00%  
Crude Fiber, Max. . . . . 0.15%  
Calcium, Min. . . . . 0.75%  
Calcium, Max. . . . . 1.25%  
Phosphorus, Min. . . . . 0.70%  
Vitamin A, Min. . . . . 35,000 IU/lb  
Vitamin D<sub>3</sub>, Min. . . . . 7,500 IU/lb  
Vitamin E, Min. . . . . 150 IU/lb

## INGREDIENTS

Dried Whey, Dried Whey Protein Concentrate, Dried Whey Product, Animal and Vegetable Fat (Preserved with BHA and BHT), Dried Skimmed Milk, Lecithin, Dicalcium Phosphate, Calcium Carbonate, Yeast Culture, Yeast Extract, Hydrolyzed Yeast, L-Lysine Monohydrochloride, DL-Methionine, Vitamin A Supplement, Vitamin D<sub>3</sub> Supplement, Vitamin E Supplement, Ascorbic Acid, Magnesium Oxide, Zinc Sulfate, Ferrous Sulfate, Niacin Supplement, Manganese Sulfate, Calcium Pantothenate, Vitamin B<sub>12</sub> Supplement, Thiamine Mononitrate, Riboflavin Supplement, Biotin, Copper Sulfate, Pyridoxine Hydrochloride, Ethylenediamine Dihydriodide, Folic Acid, Choline Chloride, Cobalt Sulfate, Menadione Sodium Bisulfite Complex (Source of Vitamin K Activity), Selenium Yeast, Sodium Silico Aluminate, Mono and Diglycerides of Edible Fats or Oils, Natural and Artificial Flavors.

SEE FEEDING DIRECTIONS ON REVERSE SIDE

Net Weight 50 lb (22.68 kg)

This product was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feed.

Manufactured for:

Eldon C. Stutsman, Inc.  
Hills, IA 52235 U.S.A.  
(319) 679-2281

MADE IN THE U.S.A.

488732

REV 1117

FI - 001

## FEEDING PROGRAM

No whole milk is required after the colostrum feeding period when calf milk replacer is used.

MIXING AND FEEDING INSTRUCTIONS	
Mix milk replacer at a rate of 8 oz of dry powder* per 2 quarts of water (110° F).	
Calves will vary in feed requirements depending on size and age. For proper medication levels (Lasalocid at 45 mg/ 100 lb body weight) - see chart. Each quart of milk replacer will contain 11.25 mg of lasalocid.	
Calf Weight (lb)	Quarts Fed per Feeding - Twice a Day
75	1-1/2
100	2
125	2-1/2

\*Dry powder should be weighed to determine recommended amount. Cup in each bag has approximate measured amounts indicated.

### FOR BEST RESULTS:

- Provide starter (grain mix) for calves when they are about 4 days of age and feed free choice up to appetite.
- Calves can be weaned when they consume 2 to 3 pounds of starter feed for 3 consecutive days. Weaning time may need to be adjusted for smaller calves.

### GOOD MANAGEMENT PRACTICES:

- Good newborn care is crucial to maximize calf survival rates. Immediate feeding of colostrum is one of the most important steps to increase the survival and health of newborn calves. Feed 2 quarts of high quality, first milking colostrum within the first 30 minutes after birth. Feed an additional 2 quarts at 6 to 12 hours of age. Continue feeding colostrum through the third day after birth.
- Begin your milk replacer program on the 4th day. Feed twice daily, at regular intervals, according to label directions. Wash equipment thoroughly after each usage.
- Have fresh, clean water available at all times.
- Do not feed hay until after weaning, when calves are consuming adequate amounts of dry starter.
- Calves should be housed in clean, dry individual pens that are well ventilated and draft free. Avoid sudden changes in temperature.

REV 1117

488732

Celmanax® is a registered trademark of Church & Dwight Co., Inc.

## FEEDING PROGRAM

No whole milk is required after the colostrum feeding period when calf milk replacer is used.

MIXING AND FEEDING INSTRUCTIONS	
Mix milk replacer at a rate of 8 oz of dry powder* per 2 quarts of water (110° F).	
Calves will vary in feed requirements depending on size and age. For proper medication levels (Lasalocid at 45 mg/ 100 lb body weight) - see chart. Each quart of milk replacer will contain 11.25 mg of lasalocid.	
Calf Weight (lb)	Quarts Fed per Feeding - Twice a Day
75	1-1/2
100	2
125	2-1/2

\*Dry powder should be weighed to determine recommended amount. Cup in each bag has approximate measured amounts indicated.

### FOR BEST RESULTS:

- Provide starter (grain mix) for calves when they are about 4 days of age and feed free choice up to appetite.
- Calves can be weaned when they consume 2 to 3 pounds of starter feed for 3 consecutive days. Weaning time may need to be adjusted for smaller calves.

### GOOD MANAGEMENT PRACTICES:

- Good newborn care is crucial to maximize calf survival rates. Immediate feeding of colostrum is one of the most important steps to increase the survival and health of newborn calves. Feed 2 quarts of high quality, first milking colostrum within the first 30 minutes after birth. Feed an additional 2 quarts at 6 to 12 hours of age. Continue feeding colostrum through the third day after birth.
- Begin your milk replacer program on the 4th day. Feed twice daily, at regular intervals, according to label directions. Wash equipment thoroughly after each usage.
- Have fresh, clean water available at all times.
- Do not feed hay until after weaning, when calves are consuming adequate amounts of dry starter.
- Calves should be housed in clean, dry individual pens that are well ventilated and draft free. Avoid sudden changes in temperature.

REV 1117

488732

Celmanax® is a registered trademark of Church & Dwight Co., Inc.

## FEEDING PROGRAM

No whole milk is required after the colostrum feeding period when calf milk replacer is used.

MIXING AND FEEDING INSTRUCTIONS	
Mix milk replacer at a rate of 8 oz of dry powder* per 2 quarts of water (110° F).	
Calves will vary in feed requirements depending on size and age. For proper medication levels (Lasalocid at 45 mg/ 100 lb body weight) - see chart. Each quart of milk replacer will contain 11.25 mg of lasalocid.	
Calf Weight (lb)	Quarts Fed per Feeding - Twice a Day
75	1-1/2
100	2
125	2-1/2

\*Dry powder should be weighed to determine recommended amount. Cup in each bag has approximate measured amounts indicated.

### FOR BEST RESULTS:

- Provide starter (grain mix) for calves when they are about 4 days of age and feed free choice up to appetite.
- Calves can be weaned when they consume 2 to 3 pounds of starter feed for 3 consecutive days. Weaning time may need to be adjusted for smaller calves.

### GOOD MANAGEMENT PRACTICES:

- Good newborn care is crucial to maximize calf survival rates. Immediate feeding of colostrum is one of the most important steps to increase the survival and health of newborn calves. Feed 2 quarts of high quality, first milking colostrum within the first 30 minutes after birth. Feed an additional 2 quarts at 6 to 12 hours of age. Continue feeding colostrum through the third day after birth.
- Begin your milk replacer program on the 4th day. Feed twice daily, at regular intervals, according to label directions. Wash equipment thoroughly after each usage.
- Have fresh, clean water available at all times.
- Do not feed hay until after weaning, when calves are consuming adequate amounts of dry starter.
- Calves should be housed in clean, dry individual pens that are well ventilated and draft free. Avoid sudden changes in temperature.

REV 1117

488732

Celmanax® is a registered trademark of Church & Dwight Co., Inc.

## FEEDING PROGRAM

No whole milk is required after the colostrum feeding period when calf milk replacer is used.

MIXING AND FEEDING INSTRUCTIONS	
Mix milk replacer at a rate of 8 oz of dry powder* per 2 quarts of water (110° F).	
Calves will vary in feed requirements depending on size and age. For proper medication levels (Lasalocid at 45 mg/ 100 lb body weight) - see chart. Each quart of milk replacer will contain 11.25 mg of lasalocid.	
Calf Weight (lb)	Quarts Fed per Feeding - Twice a Day
75	1-1/2
100	2
125	2-1/2

\*Dry powder should be weighed to determine recommended amount. Cup in each bag has approximate measured amounts indicated.

### FOR BEST RESULTS:

- Provide starter (grain mix) for calves when they are about 4 days of age and feed free choice up to appetite.
- Calves can be weaned when they consume 2 to 3 pounds of starter feed for 3 consecutive days. Weaning time may need to be adjusted for smaller calves.

### GOOD MANAGEMENT PRACTICES:

- Good newborn care is crucial to maximize calf survival rates. Immediate feeding of colostrum is one of the most important steps to increase the survival and health of newborn calves. Feed 2 quarts of high quality, first milking colostrum within the first 30 minutes after birth. Feed an additional 2 quarts at 6 to 12 hours of age. Continue feeding colostrum through the third day after birth.
- Begin your milk replacer program on the 4th day. Feed twice daily, at regular intervals, according to label directions. Wash equipment thoroughly after each usage.
- Have fresh, clean water available at all times.
- Do not feed hay until after weaning, when calves are consuming adequate amounts of dry starter.
- Calves should be housed in clean, dry individual pens that are well ventilated and draft free. Avoid sudden changes in temperature.

REV 1117

488732

Celmanax® is a registered trademark of Church & Dwight Co., Inc.