



# PERRY MILLING



# **Performance 16% Goat** Grower

### **MEDICATED**

A Complete Feed for Goats.

For the prevention of coccidiosis caused by Eimeria crandalis, Eimeria christenseni and Eimeria ninakohlyakimovae.

### **ACTIVE DRUG INGREDIENT**

Monensin (as monensin	20.0 g/ton						
GUARANTEED ANALYSIS							
Crude Protein			(Min)	16.0 %			
* (This includes not more than 1.70 % equivalent non-protein nitrogen)							
Crude Fat			(Min)	5.4 %			
Crude Fiber			(Max)	10.0 %			
Acid Detergent Fiber			(Max)	11.0 %			
Calcium	(Min)	8.0	(Max)	1.2 %			
Phosphorus			(Min)	0.4 %			
Salt	(Min)	0.4	(Max)	0.6 %			
Zinc			(Min)	80.0 ppm			
Copper			(Min)	30.0 ppm			
Selenium			(Min)	0.1 ppm			
Vitamin A			(Min)	26,000.0 IU/lb			
Vitamin E			(Min)	26.0 IU/lb			

### LIST OF INGREDIENTS

Corn, Corn Distillers Dried Grains with Solubles, Soybean Hulls, Wheat Middlings, Molasses Products, Soybean Meal, Calcium Carbonate, Soybean Oil, Ammonium Chloride, Salt, Copper Sulfate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid, Zinc Sulfate, Zinc Oxide, Ferrous Sulfate, Manganese Sulfate, Manganese Oxide, Ethylenediamine Dihydriodide, Calcium lodate, Cobalt Sulfate, Cobalt Carbonate, Sodium Selenite

### FEEDING DIRECTIONS

Feed at a rate of 3.0 to 5.5% of animal body weight to reach desired body condition. Typical consumption is 3 lbs of grain mix. Provide free choice good quality alfalfa hay and/or grass hay. Provide an abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further feeding directions consult your supplier or nutritionist.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats. CAUTION: This feed contains added copper, do not feed to sheep. CAUTION: Do not exceed 0.3 mg/kg of selenium in the complete feed.

### Manufactured By:

Perry Milling Feed & Supply LLC 102 West Front Street Perry, KS 66073 (785) 597-5122

No warranty of results is made due to differences in environment, genetics, health, and management. This product was made in a feed manufacturing facility that does not handle or store products containing animal protein prohibited in ruminant feed. g160 0pk performance 16% goat grower feed (180309) 12/16/2022

NET WT 50 LB (22.68 kg) or BULK

Code No.: G160 0PK

MEDICATED







EST. FEED & SUPPLY 1992 CONTENTS LABORET TO STATE OF THE PROPERTY LABORET TO STATE OF THE PROPERTY



# Performance 16% Goat Grower

### **MEDICATED**

A Complete Feed for Goats.

For the prevention of coccidiosis caused by Eimeria crandalis, Eimeria christenseni and Eimeria ninakohlyakimovae.

#### ACTIVE DRING INCREDIENT

ACTIVE DRUG INGREDIENT								
Monensin (as monensin	20.0 g/ton							
GUARANTEED ANALYSIS								
Crude Protein			(Min)	16.0 %				
* (This includes not more than 1.70 % equivalent non-protein nitrogen)								
Crude Fat			(Min)	5.4 %				
Crude Fiber			(Max)	10.0 %				
Acid Detergent Fiber			(Max)	11.0 %				
Calcium	(Min)	8.0	(Max)	1.2 %				
Phosphorus			(Min)	0.4 %				
Salt	(Min)	0.4	(Max)	0.6 %				
Zinc			(Min)	80.0 ppm				
Copper			(Min)	30.0 ppm				
Selenium			(Min)	0.1 ppm				
Vitamin A			(Min)	26,000.0 IU/lb				
Vitamin E			(Min)	26.0 IU/lb				

#### LIST OF INGREDIENTS

Corn, Corn Distillers Dried Grains with Solubles, Soybean Hulls, Wheat Middlings, Molasses Products, Soybean Meal, Calcium Carbonate, Soybean Oil, Ammonium Chloride, Salt, Copper Sulfate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid, Zinc Sulfate, Zinc Oxide, Ferrous Sulfate, Manganese Sulfate, Manganese Oxide, Ethylenediamine Dihydriodide, Calcium Iodate, Cobalt Sulfate, Cobalt Carbonate, Sodium Selenite

### FEEDING DIRECTIONS

Feed at a rate of 3.0 to 5.5% of animal body weight to reach desired body condition. Typical consumption is 3 lbs of grain mix. Provide free choice good quality alfalfa hay and/or grass hay. Provide an abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further feeding directions consult your supplier or nutritionist.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats. CAUTION: This feed contains added copper, do not feed to sheep. CAUTION: Do not exceed 0.3 mg/kg of selenium in the complete feed.

### Manufactured By:

Perry Milling Feed & Supply LLC 102 West Front Street Perry, KS 66073 (785) 597-5122

No warranty of results is made due to differences in environment, genetics, health, and management. This product was made in a feed manufacturing facility that does not handle or store products containing animal protein prohibited in ruminant feed. g160 0pk performance 16% goat grower feed (180309) 12/16/2022

NET WT 50 LB (22.68 kg) or BULK

Code No.: G160 0PK

MEDICATED





## PERRY MILLING





# Performance 16% Goat Grower

### **MEDICATED**

A Complete Feed for Goats.

For the prevention of coccidiosis caused by Eimeria crandalis, Eimeria christenseni and Eimeria ninakohlyakimovae.

#### **ACTIVE DRUG INGREDIENT**

Monensin (as monensin	20.0 g/ton						
GUARANTEED ANALYSIS							
Crude Protein			(Min)	16.0 %			
* (This includes not more than 1.70 % equivalent non-protein nitrogen)							
Crude Fat			(Min)	5.4 %			
Crude Fiber			(Max)	10.0 %			
Acid Detergent Fiber			(Max)	11.0 %			
Calcium	(Min)	8.0	(Max)	1.2 %			
Phosphorus			(Min)	0.4 %			
Salt	(Min)	0.4	(Max)	0.6 %			
Zinc			(Min)	80.0 ppm			
Copper			(Min)	30.0 ppm			
Selenium			(Min)	0.1 ppm			
Vitamin A			(Min)	26,000.0 IU/lb			
Vitamin E			(Min)	26.0 IU/lb			

#### LIST OF INGREDIENTS

Corn, Corn Distillers Dried Grains with Solubles, Soybean Hulls, Wheat Middlings, Molasses Products, Soybean Meal, Calcium Carbonate, Soybean Oil, Ammonium Chloride, Salt, Copper Sulfate, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Mineral Oil, Silicon Dioxide, Ethoxyquin (a preservative up to 12.0%), Butylated Hydroxyanisole (a preservative), Citric Acid (a preservative), Phosphoric Acid, Zinc Sulfate, Zinc Oxide, Ferrous Sulfate, Manganese Sulfate, Manganese Oxide, Ethylenediamine Dihydriodide, Calcium Iodate, Cobalt Sulfate, Cobalt Carbonate, Sodium Selenite.

### **FEEDING DIRECTIONS**

Feed at a rate of 3.0 to 5.5% of animal body weight to reach desired body condition. Typical consumption is 3 lbs of grain mix. Provide free choice good quality alfalfa hay and/or grass hay. Provide an abundant, fresh supply of water. Provide a separate source of salt and mineral, free choice. For further feeding directions consult your supplier or nutritionist.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats. CAUTION: This feed contains added copper, do not feed to sheep. CAUTION: Do not exceed 0.3 mg/kg of selenium in the complete feed.

### Manufactured By:

Perry Milling Feed & Supply LLC 102 West Front Street Perry, KS 66073 (785) 597-5122

No warranty of results is made due to differences in environment, genetics, health, and management. This product was made in a feed manufacturing facility that does not handle or store products containing animal protein prohibited in ruminant feed. g160 0pk performance 16% goat grower feed (180309) 12/16/2022

NET WT 50 LB (22.68 kg) or BULK

Code No.: G160 0PK

MEDICATED